

Restriction
Endonuclease



Rga I

Recognition
Sequence:

GCGAT↓CGC
CGC↑TAGCG

S

E491

200 units
5,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	75-100	50-75	10-25	25-50	100	100

55°C

80°C

Y

Ad2

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Rhizobium galegae*.

Supplied in:

20 mM Tris-HCl (pH 7.5), 250 mM NaCl, 0.1 mM EDTA,
1 mM DTT, 50% glycerol.

Reaction Conditions:

1X SE-Buffer Y. Incubate at 55° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac 66 mM KAc
10 mM MgAc 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Adv-2 DNA in 1 hour at 55° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 5-fold overdigestion with Rga I, more than 90% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 5 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.
High enzyme concentration may result in star activity.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 5 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y.

Blocked by overlapping Dam methylation (G^mATC):
GCGATCGC

Blocked by CpG methylation.