

Restriction  
Endonuclease



# Acc65 I

Recognition  
Sequence:

G↓GTACC  
CCATG↑G

S

E003

1,000 units  
20,000 u/ml

Lot:

Exp:

Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	10-25	25-50	75-100	100	10-25	100

37°C

65°C

W

λ

Dcm

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Acinetobacter calcoaceticus 65*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 50 mM KCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:

1x SE-Buffer W. Incubate at 37°C.

1X SE-Buffer W (pH 8.5 @ 25°C):

10 mM Tris-HCl    100 mM NaCl  
10 mM MgCl<sub>2</sub>    1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65 °C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of λ DNA in 1 hour at  
37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation:After 10-fold overdigestion with Acc65 I,  
> 90% of the DNA fragments can be ligated with T4  
DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg  
of DNA and 20 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of  
a single-stranded and double-stranded oligonucleotide  
was observed after incubation with 10 units of restriction  
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer W.

Blocked by overlapping Dcm-methylation (C<sup>m</sup>CWGG):  
GGTACC**CCWGG**.